

Anthropogenic impacts on the Yellow River Basin

Shuai Wang 1, Shuang Song 1,2,3, Haoyu Zhang 1, Lu Yu 1, Chentai Jiao 1, Changjia Li 1, Xutong Wu 1, Wenwu Zhao 1, Jim Best 4,5,6,7, Patrick Roberts & Bojie Fu 8 ⊠

Abstract

The Yellow River Basin supports a population of 200 million people and contains 15% of arable land in China. Water scarcity in the region is being exacerbated by climate change and human activities. In this Review, we discuss anthropogenic impacts on the hydrological cycle and sediment dynamics of the Yellow River since the 1950s. The Yellow River had one of the largest sediment loads in the world, peaking at 2.1 Gt yr⁻¹ in 1958. Such high sediment loads elevated flood risk; therefore, reservoirs, conservation and revegetation projects were implemented, reducing sediment transport by 90% since the 1980s. However, these efforts also impacted the hydrology of the Yellow River Basin, leading to an increase in evapotranspiration fluxes (1.79 mm yr⁻², 1980–2020) and reduced runoff. In addition, human water use has increased by 15.8% since the 1980s. The resulting reductions in soil water storage and intensification of the vertical water cycle foreshadow potential resource crises and will potentially lead to irreversible ecosystem degradation. Predicting the outcomes of water management policies and engineering projects is essential but highly complex owing to feedback loops and interactions between human activities and hydrological changes. Addressing these challenges, which are also faced by other arid-region rivers, will require dynamic monitoring of water storage and improved understanding of human-hydrological interactions.

¹State Key Laboratory of Earth Surface Processes and Disaster Risk Reduction, Faculty of Geographical Science, Beijing Normal University, Beijing, China. ²Department of Co-evolution of Land Use and Urbanisation, Max Planck Institute of Geoanthropology, Jena, Germany. ³Department Structural Changes of the Technosphere, Max Planck Institute of Geoanthropology, Jena, Germany. ⁴Department of Earth Science and Environmental Change, University of Illinois Urbana-Champaign, Urbana, IL, USA. ⁵Department of Geography and GIS, University of Illinois Urbana-Champaign, Urbana, IL, USA. ⁶Department of Mechanical Science and Engineering, University of Illinois Urbana-Champaign, Urbana, IL, USA. ⁷Ven Te Chow Hydrosystems Laboratory, Department of Civil and Environmental Engineering, University of Illinois Urbana-Champaign, Urbana, IL, USA. ⁸State Key Laboratory of Regional and Urban Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, China. ⊠e-mail: bfu@rcees.ac.cn

Sections

Introduction

Hydrological changes

Changing sediment load and deposition

Anthropogenic changes

Interplay of human and hydrological systems

Summary and future perspectives

Introduction

Anthropogenic pressures driven by population growth, economic development and rising water demand¹⁻³ are increasingly threatening the integrity and sustainability of river basin ecosystems⁴⁻⁶, putting the future of over 2 billion people living in drylands at risk^{7,8}. The Yellow River (also called the Huanghe) is the fifth-longest river in the world and spans arid to semi-arid regions (Supplementary ws. 1 and 2). However, the annual water discharge of the Yellow River is disproportionately lower than that of other major rivers. Anthropogenic impacts have profoundly altered – and continue to shape – the hydrological cycle, soil erosion, sediment transport and sediment deposition in the Yellow River Basin (YRB)⁹. Ecosystem degradation in the source region of the Yellow River¹⁰, severe soil erosion in the middle reaches, and depleted downstream streamflow have resulted in multiple socioeconomic stresses alongside water resources conflicts and eco-economic poverty¹¹⁻¹⁴.

The Yellow River has one of the highest sediment loads in the world ^{13,15-17}. The susceptibility of loess to erosion, combined with periodic extreme rainfall events, sparse vegetation and long-term and intensive farming activities, have resulted in severe soil erosion on the Loess Plateau in the middle reaches of the Yellow River. This soil erosion contributes to the high sediment loads, increasing the risk of flooding ^{16,18}. Since the 1980s, reservoir operation, soil conservation and large-scale reforestation have reduced sediment flux by over 90% (ref. 16). The total water storage of the Yellow River reservoirs (50 billion cubic metres) now exceeds the total annual discharge ⁹, marking the YRB as a controlled river basin where people effectively regulate its water and sediment ^{19,20}.

The YRB is facing multiple water scarcity challenges. The YRB supports 12% of the national population and 15% of arable land, with merely 2% of China's water resources 21,22. Water withdrawal has increased to 80% of the Yellow River's annual discharge, leading to perennial flow interruptions in the lower reaches that exceed the threshold for ecological collapse. Efforts to reduce soil erosion through ecological restoration have also increased water consumption¹¹, primarily driven by enhancement of evapotranspiration 23,24. In addition, runoff reduction 25 and soil moisture depletion 26 have been observed across the basin (Fig. 1). The ecological, social and economic consequences of diminishing water availability are thus becoming critical issues for the YRB^{12,14}.

In this Review, we discuss the impacts of human activities on water and sediment dynamics in the YRB. First, we outline long-term trends and the magnitude of changes in water flow and sediment load in the YRB. We then identify changes in human population, water use, land use, vegetation and economy and explore feedback loops between these changes and the basin hydrology. Finally, we discuss areas in which further research is needed to address uncertainties in human–river feedbacks and hydrological outcomes, including the feedback between precipitation and vegetation restoration, links between soil erosion and delta dynamics, and the cascading and lock-in effects of water governance.

Hydrological changes

Various data sources (Supplementary Table 1), including station observations, remote sensing data and reanalysis data, show that the components of the YRB water cycle have changed substantially since the 1950s (Fig. 1). For example, the discharge of the Yellow River has significantly decreased 9,27 at -5.78×10^8 m³ yr $^{-1}$ during 1950–2020 (P<0.01) without marked changes in precipitation. The vertical water vapour

cycle has intensified, with increased evapotranspiration but with soil moisture, groundwater and glacier storage all decreasing ^{11,27,28}. The rising human demand for water resources has driven a rapid increase in water withdrawals. Expansion in the number and area of reservoirs has also led to an increase in open water bodies, which are then exposed to evaporation ^{29,30}. Consequently, the water cycle of the YRB has exhibited a decrease in blue water (available surface water and groundwater), an increase in green water flow (evapotranspiration, especially transpiration) and a reduction in green water storage (soil moisture) ^{31,32}, as now discussed.

Precipitation

Long-term average annual precipitation in the YRB is approximately $472.69\pm94.16~\text{mm}~\text{yr}^{-1}$, decreasing from southeast to northwest and predominantly concentrated between May and September 33,34 . During 1960-2000, precipitation in the YRB decreased at $-2.21~\text{mm}~\text{yr}^{-2}$ $(P<0.05)^{35-38}$, but subsequently increased with a rate of $3.12~\text{mm}~\text{yr}^{-2}$ in 2000-2022~(P<0.05) (Fig. 2a). Precipitation thus does not display a clear statistically significant long-term trend. There is substantial spatial variability in precipitation trends across the YRB. For example, precipitation increased in the source reaches of the YRB, its upper regions and the Loess Plateau 39 between 1960 and 2020 but decreased elsewhere in the basin 35,38 (Fig. 2b). In addition, the frequency and concentration of extreme precipitation events in the YRB is increasing $^{38,40-42}$. This increase in extreme events has intensified flood risk and drought vulnerability, making the management of reservoirs and scheduling of water releases more complicated 43 .

Evapotranspiration

Between 1980 and 2020, the YRB had an annual average evapotran-spiration of approximately 370.69 \pm 55.92 mm yr $^{-1}$. Thus, over 75% of yearly precipitation evaporates 44 , which far exceeds the global average $(-60\%)^{45}$. Despite marked differences between data sources (Supplementary Table 1), all estimations consistently show an increasing trend in evapotranspiration — with up to 86% of the basin area showing a significant rise (α = 0.05) (Fig. 2c,d) whereas soil evaporation exhibits a decreasing trend at -0.42 mm yr $^{-2}$ (P<0.01) 23,46 . This increase in evapotranspiration is attributed primarily to increased temperatures and establishment of vegetation through ecological restoration 24,46 . Irrigation also contributes to the increase in evapotranspiration in the YRB, accounting for up to 44.3% of the total increase in evapotranspiration in areas with large irrigation complexes 41,44,47 .

Soil moisture and groundwater

There is substantial variability in estimates of soil moisture in the YRB^{48,49}. For instance, during 1980–2020, ERA5 data suggest a decrease in soil moisture, whereas Global Environmental Assessment Model (GLEAM) data show an increase (Fig. 2e). Observations of soil moisture show a decline from 1980, which is particularly pronounced in the middle and lower reaches of the YRB^{28,50,51} (Fig. 2f and Supplementary Table 2). This decrease in soil moisture might be attributed partly to enhanced evapotranspiration caused by increased vegetation cover 11,52.

Although available data on groundwater are limited, most results suggest that groundwater in the YRB is declining (Supplementary Table 2). This decrease is driven primarily by intensive groundwater extraction^{53,54}. Monitoring wells suggest that there is high spatial heterogeneity in groundwater changes⁵⁵; however, large-scale spatial estimations based on Gravity Recovery and Climate Experiment (GRACE) data show a continuous decline at a rate of approximately –3 mm yr⁻¹




Fig. 1| **The water cycle in the YRB.** A schematic diagram of the magnitude and trends (for the period indicated in parentheses) of components of the Yellow River Basin (YRB) water cycle where such information is available and physically meaningful, including precipitation (data are from CNO5 (ref. 197)), evapotranspiration (Global Environmental Assessment Model (GLEAM)v3.8 (ref. 198)), soil moisture (ERA5 (ref. 199)), discharge for the source area (recorded at the Tangnaihai hydrological station (Supplementary Table 1)), discharge and

sediment load in the lower reaches (recorded at the Lijin station (Supplementary Table 1)), the precipitation recycling ratio (calculated using the Lagrangian moisture tracking model data²⁰⁰), human water use (calculated based on the Yellow River Basin Water Resources Bulletin data (Supplementary Table 1)), opensurface water bodies²⁹, reservoir capacity⁹¹⁹, glacier area⁶⁵, permafrost depth⁶⁶, groundwater⁵³ and the proportion of human water withdrawals³⁵. Many of these components of the YRB water cycle have changed substantially since the 1950s.

in groundwater storage^{41,53,55,56}. The decline in groundwater and soil moisture underscores the severity of water resource shortages in the YRB.


Glaciers and open-surface water bodies

The warming climate has reduced glacier area by 21.35 km 2 (1960s–2014) and caused a decreasing trend in permafrost depth of -0.45 mm yr $^{-1}$ (1961–2017) in the source region of the YRB (Fig. 1). Approximately 35–42% of the runoff of the Yellow River originates from the


Qinghai-Tibet Plateau $^{57-61}$. Accelerated glacier retreat and permafrost melting alter the timing and magnitude of meltwater inputs, potentially causing seasonal shifts in water availability and exacerbating early spring flooding and late-season low flows $^{62-64}$. Therefore, these changes might weaken water supply capacity and threaten water availability $^{64-66}$.

The area of open-surface water bodies, including rivers, lakes, reservoirs and other terrestrial water bodies is increasing. The construction and operation of large reservoirs in the basin to support the increased proportion of discharge used by humans (from less than 20%


a Mean annual precipitation



b Spatial pattern of precipitation change




C Mean annual evapotranspiration




d Spatial pattern of evapotranspiration change




e Mean annual soil water



f Spatial pattern of soil water change



g Mean annual water discharge



h Change in water discharge




Fig. 2| **Time series and trends in water cycle components. a**, A time series of annual mean precipitation in the Yellow River Basin (YRB) (the data are from CN05 (refs. 197,201), ERA5 (ref. 199), CHM-pre 202 and CMFD 203). The red dashed and solid lines show the mean values of multiple datasets and corresponding trend lines, respectively. **b**, The spatial patterns of precipitation change between 1980 and 2022 in the YRB (the data are from CN05 (ref. 197)), the stippled area indicates regions with a statistically significant change (α = 0.05). **c**, Same as in **a** but for the mean annual evapotranspiration (the data are from the Global Environmental Assessment Model (GLEAM) 204 , ERA5 (ref. 199), MODIS16 (ref. 205) and a synthesized data package 206). **d**, Same as in **b** but for evapotranspiration (the data are from GLEAM v3.8 (ref. 198)). **e**, Same as in **a** but

for mean annual soil water (the data are from GLEAM²⁰⁴, ERA5 (ref. 199), SMAP²⁰⁷ and GLADS²⁰⁸). **f**, Same as in **b** but for soil moisture (the data are from ERA5 (ref. 199)). **g**, A time series of mean annual water discharge (dashed line) and the average trend (solid line) measured at hydrological stations at Tangnaihai (TNH), Toudaoguai (TDG), Huayuankou (HYK) and Lijin (LJ). The data are derived from the Yellow River Conservancy Commission (Supplementary Table 1). **h**, The location of the hydrological stations in **g** and the change in water discharge measured at each between 1960 and 2020. The YRB has exhibited no marked changes in precipitation patterns, significant evapotranspiration increases, soil moisture depletion and changes in discharge that differ between upstream and downstream regions.

in the 1950s to over 85% in 2010s) has led to a rapid increase in the capacity of open water bodies ²⁹. Remote sensing data show that between 1990 and 2020 the area of open-surface water bodies increased by 27.5% during floods and 58.9% during the dry season ²⁹. The five largest reservoirs along the main course of the Yellow River account for 9.48% of the total area of water bodies in the basin ^{29,35}. The total capacity of the YRB reservoirs exceeds the average annual discharge of the Yellow River ^{67,68}.

River discharge

River discharge in the YRB has decreased, owing to increased evapotranspiration and changes in land use 35,44 (Fig. 2g). Between 1950 and 2020, there was no substantial overall decline in discharge in the upper reaches and source regions of the Yellow River. Conversely, discharge in the middle and lower reaches declined at $-3.93\times10^8\,\text{m}^3\,\text{yr}^{-1}$ and $-5.78\times10^8\,\text{m}^3\,\text{yr}^{-1}$ (P<0.01), respectively. This decline has gradually stabilized since the year $2000^{9.35,69}$ (Fig. 2h). Observed changes in actual discharge are linked not only to a warmer and drier climate, but also to increased human water consumption and changes in vegetation cover $^{9.37,41}$. Hydrological data from multiple stations along the river indicate that downstream discharge is lower than upstream discharge owing to human water use 27,41 .

Vegetation restoration alters runoff by direct water consumption and through vegetation–climate feedbacks. Since the 1980s, the Yellow River has undergone extensive vegetation restoration $^{70-72}$, and it has long been believed that increased vegetation cover reduces runoff, alters the distribution of precipitation and increases evapotranspiration 73 . However, the importance of feedbacks between vegetation and the atmosphere is being increasingly recognized. Observational data and numerical simulations reveal that increased vegetation cover intensifies land–atmosphere interactions, thereby enhancing the local water cycle and increasing precipitation 74,75 . Long-term observations (1980–2020) exhibit robust increasing trends (2.76 mm yr $^{-2}$) of surface water yield over 82.3% of the Loess Plateau since revegetation commenced 39 . This seemingly counterintuitive trend highlights the importance of considering vegetation–climate feedbacks in assessing hydrological responses to large-scale vegetation changes and restoration efforts.

Changing sediment load and deposition

The Yellow River has historically been characterized by an extremely high sediment load 5,15,16 . This section outlines the temporal and spatial patterns in sediment loads and the effects of efforts to reduce the sediment load.

Temporal trends

The Yellow River used to have one of the largest sediment loads in the world⁵ (Supplementary Fig. 3). Before 1000 A.D. ^{15,16,76,77}, sediment load


in the Yellow River was at a level of approximately $0.2 \, \mathrm{Gt} \, \mathrm{yr}^{-1}$ (Fig. 3a) but then began to gradually increase owing to increasing soil erosion caused by cultivation and land use change on the Loess Plateau. In the 1950s, sediment concentrations in the Yellow River surpassed $40 \, \mathrm{kg} \, \mathrm{m}^{-3}$ (refs. 78,79) (Fig. 3b) with an average annual sediment load of $1.6 \, \mathrm{Gt} \, \mathrm{yr}^{-1}$. The sediment load peaked in 1958 at $2.1 \, \mathrm{Gt}$, far exceeding the natural baseline conditions 15,76,77,80 . The average sediment load of the Yellow River during $1961-1990 \, \mathrm{was} \, 0.883 \, \mathrm{Gt} \, \mathrm{yr}^{-1}$, accounting for 5-10% of global sediment flux into the oceans $^{5,81-83}$.

The once characteristically high sediment load of the Yellow River has diminished substantially owing to anthropogenic changes, including vegetation restoration, landscape engineering and the construction of reservoirs and check dams¹⁶. The sediment load of the Yellow River declined at an average rate of -0.024 Gt yr⁻² during the second half of the twentieth century. The average annual sediment load after 2000 is 0.148 Gt, only 10% of that observed in the 1950s. In addition, the sediment concentration decreased by 75% after 2000 (Fig. 3a,b).


Spatial heterogeneity

Large quantities of sediment are transported from the Loess Plateau into the Yellow River^{16,17}, which accumulate mostly in the lower reaches of the river and its estuary. Although over 60% of the Yellow River's water discharge originates from the YRB above Lanzhou⁶⁰, the sediment load at Lanzhou is merely 8% of that from the lower reach (Fig. 3c). The Loess Plateau in the middle reach contributes 90% of the sediment load entering the Yellow River^{16,70,83,84}. In the lower reach of the Yellow River, the channel gradient decreases, leading to substantial sediment deposition⁸⁵. Therefore, levees have been constructed in the lower reaches to control floods¹⁷. This intense sediment deposition has elevated the riverbed by over 10 m above the adjacent alluvial plain 18,86, forming what is commonly known as a 'hanging river' constrained within levees¹⁸ (Supplementary Fig. 1). This phenomenon has led to frequent flooding, breaching and diversion events on the downstream Yellow River floodplain 18,76,87,88. Moreover, the river channel continues to meander and silt up between the levees, with breaching of the elevated river further threatening the floodplain within, and outside, the levees⁸⁹⁻⁹¹.


Sediment deposition in the lower reaches of the Yellow River has also led to the formation of alluvial plains and deltas. The modern Yellow River Delta, which began forming in 1855 following diversion of the Yellow River to its current estuary, has created approximately $2.7\times10^3\,\mathrm{km^2}$ of new land 92 . During the last 6,000 years, sediment deposition has pushed the coastline 60 km into the Bohai Sea 93 . It is also estimated that over 530 Gt of sediment have been deposited in the alluvial plains and deltas since 1128 A.D. 77 . This deltaic expansion has created land resources and formed ecologically valuable wetlands with high



b Sediment concentrations



C Spatial pattern in water and sediment discharge



d Responses to temporal variations in sediment flux




Fig. 3 | **Spatial and temporal patterns of sediment transport. a**, A time series of sediment load in the lower reaches of the Yellow River throughout history (yellow solid line highlights the trends; the data are from refs. 5,77) and from the 1950s to the present day (the solid line and shaded region indicate the linear regression and 95% confidence interval, respectively, with the solid line (right) indicating the periodic mean of the sediment load) in the context of key events. **b**, A box and whisker plot of changes in sediment concentration in the lower reaches of the Yellow River (centre line, the top and bottom of the box and whisker extent indicate the average value, 75% and 25% quantiles and maximum and minimum value in each period, respectively). **c**, Variation and spatial pattern in water (left side of inset plots) and sediment discharge (right side of inset plots) across the source (I), upper (II, III), middle (IV, V) and lower reaches (VI)

of the Yellow River Basin (YRB). **d**, Temporal response of the bankfull discharge, (blue bars, the data are derived from ref. 98 and the Yellow River Conservancy Commission), delta area (yellow dots, the data are from refs. 95,103,104,209,210) and channel migration rate (blue dots) of the lower Yellow River in response to temporal variations in sediment flux (colour bar (top), the data are from ref. 89). The blue and yellow solid lines and shaded areas show the regression line of the generalized linear model and the 95% confidence interval for the channel migration rate and delta area variation, respectively. The data are sourced from the Yellow River Conservancy Commission (Supplementary Table 1) unless otherwise stated. The once-heightened sediment load of the Yellow River has decreased spatially and temporally, leading to changes in the downstream channel and delta.

biodiversity, various wildlife habitats and important over-wintering and breeding sites for migrating birds 94,95 . However, the stability of the Yellow River Delta is heavily influenced by changes in the dynamics of the sediment load 95 .

Responses to decreased sediment

Decreased sediment flux, as well as active reservoir regulation, have mitigated the problem of sediment deposition in the lower reaches of the Yellow River, Upstream regulation, including landscape engineering, ecological restoration and reservoir operation in the late twentieth century^{89,96}, led to a continuous decrease in the sediment load, reducing the deposition rate in the lower reaches⁸⁶. In 2000, the sediment load entering the lower reaches of the Yellow River dropped below 0.1 Gt yr⁻¹, leading to a transition from a state of deposition to net erosion⁸⁶. Furthermore, the Water-Sediment Regulation Scheme implemented after 2002 has accelerated the incision and widening of the channel^{86,96}. Consequently, the elevation of the riverbed has decreased by up to 3 m (refs. 96,97), the bankfull discharge (the maximum discharge that can be held within the channel) of the lower Yellow River increased by 200% (ref. 98), and the river channel has stabilized⁸⁹ (Fig. 3d and Supplementary Fig. 4). These changes have benefited flood control in the lower reaches of the Yellow River, enabling further agricultural development on the floodplain⁸⁹. Whether this reduced sediment load can be sustained remains a critical challenge for communities within the YRB.

Nevertheless, the evolution of reservoirs and the riverbed in the lower reaches has posed challenges to the sustainability of strategies for the mitigation of sediment deposition and flooding. For example, the Sanmenxia and Xiaolangdi reservoirs, which were built to control water and sediment flux in the lower reaches of the Yellow River in 1961 and 1999, respectively, have undergone severe siltation 99,100. Consequently, their storage capacity and the flushing efficiency in the lower reaches have reduced 86,99,101. In addition, channel erosion downstream of the dams has led to riverbed incision and coarsening of the bed sediment, because fine sediment is preferentially entrained in water flow 99,100. This bed coarsening makes the riverbed more difficult to erode 99. Furthermore, a coarsened riverbed leads to the formation of larger sand dunes in the channel, which increases flow drag. Such extra drag has been proposed to exacerbate the risk of flooding during moderate and large floods 97,102.

Reductions in sedimentation also have negative impacts on the Yellow River Delta, with delta expansion slowing since the $1990s^{79,95,103,104}$ (Fig. 3d and Supplementary Fig. 4). Despite delta growth continuing near the estuary, coastal erosion has occurred further away from the estuary ^{79,103}, leading to concerns over potential ecological degradation resulting from delta shrinkage ^{79,104,105}. In addition, the delta is vulnerable

to the effects of climate change such as sea level rise¹⁰⁶; therefore, approaches to sediment regulation in the YRB must balance the need for flood protection in the lower reaches with the sustainability of the delta.

Anthropogenic changes


Population growth, urbanization and human activities have shaped the socio-ecological landscape of the YRB since the 1950s. This section explores the complex human dimensions driving changes within the YRB, including demographic shifts, patterns of water use, land use transformations and socioeconomic developments.

Changing population distribution

Population expansion and increased demands for food and energy have intensified water stresses in the YRB. The human population in the YRB rose from 10 million in the seventh century to 100 million in 1950, 146 million in 1980 and 216 million in 2020 (Fig. 4a, Supplementary Table 1). Urbanization and population aging have led to shifts in lifestyle, consumption and production patterns, altering water use in various sectors and producing other environmental impacts¹⁰⁷ (Fig. 4b). Rapid industrialization and urbanization correlate with population increase over the same periods¹⁰⁸ (Fig. 4a). Despite the continuous population growth from 1980 to 2020, the per capita water withdrawal in the middle and lower reaches of the YRB declined from 396 m³ vr⁻¹in 1980 to 310 m³ yr⁻¹ in 2020, partly owing to stricter water resource management and transformations in production patterns. These changes highlight a pronounced spatial mismatch between water resource endowments and population distributions (Fig. 4). Moreover, water scarcity is further intensified by the prevalent migration of people towards economically developed downstream regions that are already experiencing water stress 109-1111.

Changes in water use

Water use in the YRB faces substantial pressures related to population growth, urbanization and economic activities. Water demand has increased by 15.8% since the 1980s (Fig. 4c). Between 1980 and 1997, water use increased at a rate of 1.02 km³ yr², mainly for irrigation, leading to a 63.7% (ref. 112, during 1982–2014) decrease in the discharge of the Yellow River $^{112-115}$. Industrial water use, the second largest sector, increased by 17.43% between 1980 and 2020. However, the rate of increase of industrial water use began to decline after 2000, changing from 0.28 km³ yr² before 1997 to -0.02 km³ yr² after 2000. Domestic water use, corresponding to water withdrawal for urban and rural household use and service activities, increased steadily at a rate of 0.17 km³ yr² from 1980 to 2020, owing to population growth,



C Human water use




Fig. 4 | Population and human water use. a, Changes in the rural, urban and total population in the Yellow River Basin (YRB) (the data are from City Statistical Yearbook and the National Census of China) (Supplementary Table 1). b, The distribution of the YRB population in 1980 and 2020 (the data are from City Statistical Yearbook and the National Census of China) (Supplementary Table 1). c, The contribution of agricultural, industrial, domestic and ecology sectors to human water use in the YRB (the data are from Provincial Water

Resource Bulletins (2000–2020); Nation Long-term Water Use Dataset of China (1965–2013) 211 (Supplementary Table 1). **d**, Spatial distribution of per capita water withdrawal in the YRB in 1980 and 2020 (the data are calculated from human water use data and population data) (Supplementary Table 1). The changes in the population of the YRB and economic importance of each industry between 1980 and 2020 have driven changes in water use across multiple sectors.

d Per capita water withdrawal in the YRB

especially in urban areas. Consequently, the contribution of domestic water use to total water use increased from 6.5% in 1980 to 14.3% in 2020. Ecological water use, referring to the water used to maintain ecosystems (including sediment flushing, planting trees and grass, recharging rivers and restoring dried-up regions), has expanded rapidly since the 2000s (0.31 km³ yr⁻²), owing to efforts to introduce policies to support sustainability of the river ecosystem 116 .

Although water management practices have decreased per capita water withdrawal since the 1980s, there is still a substantial disparity between water availability and demand. Severe drying-up of the river occurred during 1980s–2000s; consequently, approaches such as the Water Allocation Scheme (1998)¹¹⁷ and the construction of the Xiaolangdi Reservoir in 1999, were proposed to mitigate water scarcity^{12,118,119}. These approaches led to a downward trend in

water use $(-0.715\,\mathrm{km^3\,yr^{-2}})$ before 2000. This trend stabilized after 2004 $(-0.02\,\mathrm{km^3\,yr^{-2}})$, being balanced by high human demand. The structure of water use has changed concurrently in response to these economic and social demand shifts. For example, the contribution of agricultural water use decreased from 81.44% to 63.75% between 1980 and 2020 but remains much higher than other sectors (Fig. 4c and Supplementary Fig. 5). Despite these measures, human water withdrawal still reached 84% of the available surface water resources in 2020, breaching the 40% water scarcity threshold 120 . Per capita water availability was still only 473 m³ yr $^{-1}$ (ref. 121) until 2020, representing -23% of the national average in China, and falling well below the global threshold for severe water scarcity (500 m³ yr $^{-1}$) (Fig. 4d).


Land use and vegetation changes

Cultivation, over-grazing and human interventions have caused ecological degradation in the YRB. This degradation started in the

eleventh century, accelerated in the eighteenth century 18,40,70,122 and persisted until the $1990s^{18,123}$. By 1850, anthropogenic land use exceeded natural land cover in 43% of the Loess Plateau (Fig. 5a). From 1950 to 1960, the extent of cropland on the Loess Plateau increased by 5×10^6 ha, further reducing natural vegetation coverage 70,124 . In the source region, grassland shrinkage, degradation, and desertification have been continuous since the $1970s^{10,125}$; consequently, vegetation cover decreased in 67% of the source region between 1982 and 2001^{126} . This severe degradation could lead to increased soil erosion, increased sediment flux into the Yellow River and, consequently, intensified channel diversion 127 .

In the frozen regions of the YRB, anthropogenic warming will alter sediment flux and vegetation growth. Enhanced melting of permafrost areas in the source regions of the Yellow River on the Tibetan Plateau has increased sediment flux and channel mobility¹²⁷. Conversely, the effect of increasing temperatures in warmer areas with seasonally

a Anthropogenic land use exceeding 50%



C Land use change (1990-2020)






Fig. 5 | **Land use, vegetation and economic changes. a**, Spatial distribution of the years when the proportion of anthropogenic land use exceeded 50% in the Yellow River Basin (YRB) (the data are from HYDE 3.3 (ref. 212)). **b**, Trend in leaf area index (LAI) between 1982 and 2020 (data from GIMMS LAI4g²¹³). **c**, A chord diagram of land use change in the YRB between 1990 and 2020 (the data are from CLCD²¹⁴, the thickness of the chords represents the area of land use change with

b Leaf area index trend (1982-2020)



d Gross value added in the YRB (1980-2020)



the chord colour representing the new land use). \mathbf{d} , The total gross value added (GVA) (solid line) and contribution of agriculture, industry and service to the GVA (coloured bars) in the YRB (1980–2020) (the data are from City Statistical Yearbook and the National Census of China) (Supplementary Table 1). Since the 1980s, the YRB has experienced rapid economic growth and ecological restoration efforts have increased vegetation coverage. CNY, Chinese Yuan.

frozen ground is modulated by enhanced vegetation growth, which stabilizes river banks and reduces channel migration¹²⁷.

Various landscape engineering approaches have been implemented to control soil erosion and reduce sediment flux into the Yellow River. The construction of check dams, which intercept sediment on hillsides or in valleys, was promoted and implemented in the 1950s; by 2011, there were over 58,000 check dams in the YRB^{128,129}. Moreover, terrace farming was promoted in the middle reaches of the Yellow River 16,70,130 with the extent of terraces reaching $6.46\times10^4 \,\mathrm{km^2}$ by 2022 (ref. 131). These measures reduced sediment flux into the Yellow River in addition to ensuring that agricultural productivity was maintained to sustain a growing population 70,129,132 . However, there is a limit on the ability of landscape engineering to reduce sediment yield 16,128 . For instance, check dams can silt up and lose their retention capacity within 10 years owing to high sediment production 128 .

Various ecological and conservation projects have been implemented in the YRB, leading to increased vegetation coverage, which mitigates soil erosion. Such projects began in the 1980s and emerged rapidly throughout the late twentieth century, including the Three North Shelterbelt Forest Program (1978)^{8,123,133}, which promoted afforestation, the Natural Forest Protection Program (1998)8,123,134,135, which aimed to prevent deforestation, and the Grain for Green Program (1999)^{8,123,135,136}, which supported the conversion of cropland to forest or grassland. These projects have enhanced forest and grassland vegetation substantially in the YRB^{8,123,135,137} (Fig. 5b,c). Between 1982 and 2020, 71% of the YRB experienced a significant (P < 0.05) increase in leaf area index (Fig. 5b). In addition, the area of grassland and forest increased by 4.29×10^3 km² (1.0%) and 1.69×10^4 km² (22.8%), respectively, between 1990 and 2020 (Fig. 5c), and the fractional vegetation cover increased by 17% between the years 1999 and 2018¹³⁸. The normalized difference vegetation index increased in over 60% of the area in the source region during the 2000s, as barren land was converted to grassland^{139,140}. In addition, in the Loess Plateau, vegetation coverage increased from 31.6% in 1999 to 59.6% in 2013, reaching 65% in 2021 owing to the conversion of cropland to grasslands and forests $^{11,16,141-143}$ (Supplementary Fig. 2).

Economics and policies

Since 1980, the YRB has experienced rapid economic growth, placing increasing demands on water resources. The gross value added (GVA) of the YRB increased by an average of 1.78% per year between 1980 and 2020 (Fig. 5d). In this period, the contribution of agriculture to the total GVA in the YRB decreased from 24.5% to 7.5%, the share of service GVA rose from 27.6% to 51.4%, and the industrial share decreased from 47.9% to 41.3% (Fig. 5d and Supplementary Fig. 5). Along with these changes, the efficiency of industrial water use increased 144,145, high water-consuming and energy-consuming industries were replaced 146, and infrastructure was enhanced to promote water-saving technology and establish standard water quotas across sectors 147.

As water resources become the primary factor restricting economic growth, water-related policies have been proposed to govern the YRB. Before the 1980s, initiatives concentrated primarily on the use of large-scale engineering projects, such as reservoirs and levees, to meet the increasing demands for agricultural, industrial and domestic water and to control floods. For example, the Sanmenxia and Liujiaxia Dams were constructed to control floods and generate hydropower. Since 1980, the focus has shifted to comprehensive institutional governance¹⁹ (Supplementary Fig. 6). For instance, the Water Allocation Scheme (1987)¹⁴⁸ was the first unified plan to allocate Yellow River water across

provinces, laying the foundation for coordinated basin management. Although the goal of halting growth in water usage was not achieved in the first decade, the withdrawal of water was strictly regulated as the administrative authority of the Yellow River Conservancy Commission expanded. In 2010, the Yellow River Conservancy Commission won the Lee Kuan Yew Water Prize¹⁴⁹ for its complex management system^{9,20}, which prevented the Yellow River from drying up. The construction of the Xiaolangdi Reservoir in 1999 strengthened integrated water management through its multifunction purpose of flood control, water and sediment regulation and maintaining ecological basic flows^{105,150,151}.

After the peak in water use in 1997, enhanced water quota policies, including the Water Allocation Scheme (1998)¹¹⁷, were introduced to further strengthen unified water management from the supply side. Policies to transfer water rights and reform agricultural water prices were also implemented from the demand side^{152,153} (Supplementary Fig. 6). The aim was to give agrarian water rights to the industrial sector in exchange for fees to develop water infrastructure¹⁵⁴. From the 2010s to the 2020s, several technique-guided policies, such as Integrated Fertilizers and Water Use Management in Agriculture¹⁵⁵, were released to promote water-saving irrigation technologies (Supplementary Fig. 6). With additional further infrastructure and policies, such as Water Diversion from South to North¹⁵⁶, Strict Water Management¹⁵⁷ and the Yellow River Protection Law (2022)¹⁵⁸, efficient use of water resources has become a performance indicator for local administrative officials.

Interplay of human and hydrological systems


Anthropogenic impacts have reshaped the hydrological dynamics of the YRB and, over time, led to further feedbacks for society. Population growth and agricultural expansion have affected the YRB for thousands of years, resulting in the removal of vegetation, severe soil erosion, and increased sediment loads¹⁷. Since the 1960s, anthropogenic activities have intensified owing to a growing population and food demands. Modernization and industrialization have stimulated engineering projects, such as reservoir construction, which alter water flow and sediment transport. Meanwhile, ecological restoration projects have altered vegetation patterns and reduced water and sediment loads. This section examines the feedback mechanisms between anthropogenic interventions and hydrological changes, identifying four hypothetical feedback loops (Fig. 6) and suggesting potential future changes. Detailed institutional and engineering solutions, plans and future potential changes are given in Supplementary Fig. 6 and Supplementary Table 3.

Feedback loops

Challenges have emerged from the ongoing intertwined interactions between humans and water, with uncertainties arising from institutional, social and technical transitions and climatic changes ^{159–161}. Over different periods, the strength and influence of these drivers have varied substantially, leading to complex feedback loops that govern the interactions between human activities and the natural environment. These shifts have caused ecological and hydrological transformations, which have often reinforced each other and complicated efforts to balance environmental protection with economic and social development in the YRB.

The supply expansion feedback loop during 1960–1980 was characterised by expanding water supply programmes to meet social and economic development demands, which reduced water discharge (Fig. 6a). Rapid socioeconomic development (for example, the GVA) in this arid and semi-arid region relies heavily on water resources 162.

a Supply expansion loop (1960s-1980s)



b Resource restriction loop (1980s-2000s)



C Complex governance loop (2000s-2020s)



d Future resilience loop (future)




Fig. 6 | Feedback loops for anthropogenic impacts on the Yellow River.

 $\label{eq:approx} \textbf{a}, Supply expansion loop (1960s-1980s), characterized by the growth of water supply programmes to meet the demands of social and economic development. The coloured boxes indicate dominant loops. <math display="block">\textbf{b}, A \ resource \ restriction \ loop (1980s-2000s) \ involving \ strategies to \ restrict \ water \ demand \ and \ the initial implementation of ecological \ restoration \ projects. \\ \textbf{c}, A \ complex \ governance} \ loop (2000s-2020 \ s) \ in \ which \ institutional \ and \ engineering \ solutions \ increased \ water \ use \ efficiency, \ enhanced \ vegetation \ and \ reduced \ sediment \ yield. \\ \textbf{d}, A \ future \ resilience \ loop \ characterized \ by \ uncertainties \ from \ climate \ change \ and \ socioeconomic \ development. \ The \ four \ hypothetical \ feedback \ loops \ represent \ dynamic \ interactions \ between \ human \ and \ water \ system \ components \ across \ different \ periods.$

Therefore, engineering infrastructure, such as diversion projects, were developed to boost water supply. This water supply supported further economic activities, creating an accelerating feedback loop. The increasing demand, driven by engineering-intensive development, led to the appropriation of water resources from external regions and other stakeholders within the basin 163,164. The lack of adequate policies and institutional constraints led to further scrambles for water allocation and drying-up crises for streamflow 165. Moreover, urbanization and land development restrict vegetation cover, further impacting discharge and sediment flux 19,124.

Between 1980 and 2000, water scarcity became a key driver of institutional shifts in the YRB leading to strategies to restrict water demand²⁰. Population growth, economic development and urbanization continued to drive demand (for example, food and energy). Therefore, water scarcity curbed uncontrolled exploitation through institutional reforms and monetary policies. However, mismatched institutional structures, under which the provinces engage in uncoordinated competition for scarce water owing to economic growth pressures, led to intensified exploitation of water resources, fuelling disputes over the fairness of governance outcomes^{166,167} (Fig. 6b). In the late 1990s, ecological restoration efforts began to emerge to mitigate some of the adverse effects (such as, drying up¹¹³, wetland shrinkage⁹⁴ and increasing soil salinization¹⁶⁸) of past practices¹⁶⁹.

A complex governance loop emerged throughout 2000–2020 in which water use efficiency increased via demand-side and supply-side management 118 and strategies such as water diversions and ecological projects were introduced to enhance vegetation and reduce sediment yield 16. Increased diversity in policy and governance enhanced water use efficiency 118, reducing overall water consumption (Fig. 6c). Meanwhile, ecological conservation and restoration efforts began to control soil erosion more effectively with unified regulation also enhancing the management of sediment deposition. Large-scale vegetation conservation and landscape engineering projects were also implemented to control severe soil erosion in the YRB.

However, vegetation restoration projects also altered hydrological processes (Fig. 6c). For example, the rapid increase in vegetation has increased evapotranspiration, reduced runoff and discharge, and caused soil moisture depletion^{26,50,170}. By 2021, vegetation in the eroded sediment areas of the middle reaches of the Yellow River had nearly recovered to its ecological threshold level (65%)¹⁴². Downstream water discharge dropped by 38% from 1971–1990 to 1991–2016, with vegetation increase responsible for up to 26% of this decline^{16,25,171}. The depletion of soil moisture and terrestrial water storage has raised concerns about hydrological sustainability in the YRB. In 2008, the maximum depth of the dried soil layer reached 5 m, primarily in the

forest regions^{26,50,172,173}. This depletion limits the future effects of ecological restoration and threatens water resource availability in the middle and lower reaches of the Yellow River^{11,174,175}. Therefore, instead of extensive afforestation, mixtures of trees, shrubs and herbaceous plants adapted to local conditions should be prioritized to balance hydrological sustainability and ecological effectiveness¹³. It is also possible that increased vegetation could enhance local water cycles and increase precipitation; however, the degree and importance of this feedback are uncertain^{39,74,75}.

Future changes

Climate change is expected to alter future water availability (Fig. 6d). Engineering and institutional approaches to address soil erosion and excessive water withdrawals have curbed the overall deterioration in water resources. However, dependence on more extensive human interventions has resulted in a legacy of maximizing water resource utilization¹⁹. The increasing frequency and severity of extreme events under the impacts of climate change might diminish the stability of water supply and challenge this water management system, which was optimized under previous human-river relationships. Glacial melting 176 will also lead to the loss of stored water. Future water resources in the YRB are projected to increase by 0.218 ± 0.201 km³ yr⁻¹ under various climate change scenarios and models (Supplementary Fig. 6). These perceived improvements in water availability could lead to increased dependence on external factors, such as increasing precipitation. Underestimating the risks associated with water crises could weaken the motivation for proactive management and policy interventions in the short term. Given that institutional reforms often exhibit inherent delays, there might be further water depletion in the long term¹⁷⁷, increasing the vulnerability of the system.

The compound impacts of reduced cropland area, population decrease¹⁷⁸ and improvements in the efficiency of water use are projected to mitigate human water demand in the late twenty-first century (Fig. 6d). Models incorporating fertility policies, well-being and economic growth project that the population of the YRB will peak around 2020–2030 (Supplementary Fig. 6). However, an ageing population might lead to more uncertain behaviours in water use (Supplementary Fig. 5). Ongoing demographic change, production shifts and climate change are likely to cause a decline in cropland area until the 2100s, leading to reduced agricultural water demand. In addition, socioeconomic and technological transitions have led to improvements in the efficiency of water use. Human water demand, modelled by integrating complex system transitions – including future population changes, land use changes, economic cycle fluctuations¹⁷⁹, technological changes and shifts in human lifestyle 180 – will potentially reach a peak of 74 km³ yr⁻¹ in the mid-twenty-first century, still exceeding the $2020 \text{ level of } 67 \text{ km}^3 \text{ yr}^{-1}$.

Other projected changes, such as urbanization, forest and grass-land expansion and increased ecological water use, could increase water demand for humans and the broader ecosystem. Urbanization is expected to continue to increase and is projected to reach 75–90% by the 2100s (Supplementary Fig. 6), accelerating the increase in domestic, food and industrial water use. The expansion of forests and grasslands remains possible under different scenarios, driven by ongoing ecological restoration efforts and climate change (Supplementary Fig. 6). This potential increase in vegetation could pose risks to domestic, industrial, and irrigated water availability. Moreover, the share of ecological water use (for example, to supplement wetland, groundwater and urban greenspace) is predicted to grow and compete

with human water use for social and economic activities 12 and threaten food production 181 . High water stress in the YRB will persist unless there is a shift in the demand 109,181 .

Complex anthropogenic–hydrological feedbacks must be considered when negotiating the conflict between human demands and water. Approaches such as landscape engineering, reservoir operation and ecological restoration have been used to address the problem of sedimentation in the YRB¹⁶. However, inertia in resource allocation might result in diminishing marginal returns and a reduced ability to respond to emerging risks. Challenges such as reaching engineering saturation, soil moisture depletion and limitations on water resources can seriously compromise the efficiency of these sediment reduction strategies^{11,128,174}. Therefore, it is important to design feedback loops to be more adaptive when facing such uncertain and complex risks.

Summary and future perspectives

Despite being historically the world's most sediment-laden major river, the YRB is now experiencing a simultaneous decline in water discharge, soil moisture and groundwater alongside rising human water consumption and evapotranspiration. Consequently, the overall flux of the water cycle is intensifying, whereas water resource storage is diminishing. Since the mid-twentieth century, large-scale engineering works and policy reforms have reduced sediment loads. Compounded by drivers such as land use transformation, escalating urbanization, and evolving water usage patterns, these projects have also altered eco-hydrological processes. The complex interplay of hydrological change, human intervention and institutional policies in the YRB exemplifies a new, multifaceted era in arid river basin management: this era is characterized by short-term water supply successes but heightened long-term risks, such as groundwater depletion and mounting agricultural dependence on irrigation. These risks foreshadow a potential water crisis that, when combined with future climate pressures, underscores the need for transformative and adaptive water governance in the YRB.

Several unresolved challenges remain for the YRB that require further hydrological research, and are issues shared by major river basins globally. First, the complexities surrounding how vegetation restoration regulates river systems, especially its feedback with precipitation, remain poorly understood but could have important implications for arid rivers undergoing restoration or deforestation, such as the Tarim¹⁷⁶, Congo watershed¹⁸² and Amazon rivers^{183,184}. Second, managing the delicate balance between controlling soil erosion and safeguarding increasingly urbanized delta ecosystems introduces scientific and policy dilemmas, as seen in major deltas 185, such as those of the Mekong 186, Rhine 187 and Yangtze 188 rivers. Third, the unintended consequences of water management policies 167 – such as shifting stress from surface to groundwater resources - raise questions concerning the sustainability of interventions across transboundary rivers¹⁸ including the Colorado, Murray and Orange rivers¹⁹⁰. Thus, further research on the YRB is important for investigating globally relevant eco-hydrological challenges and developing innovative, transferable solutions.

Comprehensive monitoring is needed to better understand arid-region river systems such as the YRB 3,8,191 . Monitoring in the YRB focuses primarily on flux measurements, such as water and sediment discharge, and evapotranspiration for water resource management. There is a shortage of dynamic monitoring data of water storage in groundwater, soil moisture and glaciers. Measurements of these water storage components rely on coarse estimations using large-scale models or remote sensing methods. The long-term decline in water storage

across the YRB could lead to water resource crises that might be overlooked by traditional monitoring flux metrics^{192,193}. Thus, comprehensive monitoring networks are needed to capture dynamic variables of the water cycle, develop a digital twin of the river system based on real-world data and consider the economic and technical feasibility of different policies and forecasts¹⁹⁴. Such monitoring and simulation technologies have been widely used in water resource management in urban areas to improve water use efficiency. However, data are still lacking in sparsely populated, underdeveloped regions. Addressing these data limitations will enable integrated modelling of human-water systems and water management within the river basin¹⁹⁵.

Adaptive capacity in regional water management practices is necessary to bridge the gap between science and policy, facilitating responses to immediate challenges and long-term uncertainties in a changing climate and socioeconomic context. The unexpected outcomes of water management policies, such as the water quota policy²⁰, reveal the complexity of river basins. Theoretical and practical gaps in the pliability of scientific research and policy-making can lead to long-term legacy effects, despite achieving success in addressing immediate challenges. The lag between intervention and impact can span decades, resulting in inequality in access to water resources among various stakeholders across different periods and regions². Efforts to enhance adaptive capacity must balance the competing objectives of economic development, ecological preservation and social equity¹. For instance, future work should aim to explore the carrying capacity of water resources; reducing human interference; prioritizing nature-based solutions; the benefits of adaptive policies such as water rights trade, adapting forestry, shrubs and grassland to environmental conditions; and reforming rigid institutions^{1,196}.

It is essential that scientific knowledge from different regions and cooperation across areas is used to achieve adaptive water governance¹⁸⁹. Therefore, sustained investment in interdisciplinary collaborative platforms, including funding, the development of young scientists, international partnership networks, and standardized assessment indicators is needed to integrate the strengths of all parties and bridge the knowledge gap in sustainable river management.

Data availability

Supplementary Table 1 outlines details of the datasets used in Figs. 1-4.

Published online: 16 September 2025

References

- Basheer, M. et al. Cooperative adaptive management of the Nile river with climate and socio-economic uncertainties. Nat. Clim. Change 13, 48–57 (2023).
- Zhao, G. et al. Decoupling of surface water storage from precipitation in global drylands due to anthropogenic activity. Nat. Water https://doi.org/10.1038/s44221-024-00367-7 (2025).
- Abel, C. et al. The human-environment nexus and vegetation-rainfall sensitivity in tropical drylands. Nat. Sustain. 4, 25–U150 (2021).
- Vörösmarty, C. J. et al. Global threats to human water security and river biodiversity. Nature 467, 555–561 (2010).
- 5. Best, J. Anthropogenic stresses on the world's big rivers. *Nat. Geosci.* **12**, 7–21 (2019).
- Best, J. & Darby, S. E. The pace of human-induced change in large rivers: stresses, resilience, and vulnerability to extreme events. One Earth 2, 510–514 (2020).
- Varis, O., Taka, M. & Kummu, M. The Planet's stressed river basins: too much pressure or too little adaptive capacity? Earths Future 7, 1118–1135 (2019).
- Li, C. et al. Drivers and impacts of changes in China's drylands. Nat. Rev. Earth Environ. 2, 858–873 (2021).
- Wang, Y., Zhao, W., Wang, S., Feng, X. & Liu, Y. Yellow river water rebalanced by human regulation. Sci. Rep. 9,1–10 (2019).
- Li, X.-L. et al. Rangeland degradation on the Qinghai-Tibet plateau: implications for rehabilitation. Land Degrad. Dev. 24, 72–80 (2013).
- Feng, X. et al. Revegetation in China's Loess Plateau is approaching sustainable water resource limits. Nat. Clim. Change 6, 1019-1022 (2016).

- Cai, X. & Rosegrant, M. Optional water development strategies for the Yellow River Basin: Balancing agricultural and ecological water demands. Water Resour. Res. 40, W08S04 (2004).
- Fu, B. et al. Hydrogeomorphic ecosystem responses to natural and anthropogenic changes in the Loess Plateau of China. Annu. Rev. Earth Planet. Sci. 45, 223–243 (2017).
- Wang, Y., Song, J., Zhang, X., Sun, H. & Bai, H. Coupling coordination evaluation of water-energy-food and poverty in the Yellow River Basin, China. J. Hydrol. 614, 128461 (2022).
- Wang, H. et al. Stepwise decreases of the Huanghe (Yellow River) sediment load (1950–2005): impacts of climate change and human activities. Glob. Planet. Change 57, 331–354 (2007).
- Wang, S. et al. Reduced sediment transport in the Yellow river due to anthropogenic changes. Nat. Geosci. 9, 38–41 (2016).
- Mostern, R. & Horne, R. M. The Yellow River: a Natural and Unnatural History (Yale Univ. Press, 2021); https://doi.org/10.2307/j.ctv1vbd1d8.
- Chen, Y., Syvitski, J. P. M., Gao, S., Overeem, I. & Kettner, A. J. Socio-economic impacts on flooding: a 4000-year history of the Yellow river, China. AMBIO 41, 682–698 (2012).
- Song, S. et al. Identifying regime transitions for water governance in the Yellow River Basin. China. Water Resour. Res. 59, e2022WR033819 (2023).
- Song, S. et al. Quantifying the effects of institutional shifts on water governance in the Yellow River Basin: a social-ecological system perspective. J. Hydrol. 629, 130638 (2024).
- Xu, Z., Takeuchi, K., Ishidaira, H. & Liu, C. An overview of water resources in the Yellow River Basin. Water Int. 30, 225–238 (2005).
- Fu, B. et al. Mechanisms of human-natural system coupling and optimization of the Yellow River Basin. Bull. Natl Nat. Sci. Found. China 35, 504–509 (2021).
- 23. Jian, S., Wang, A., Hu, C. & Yan, D. Effect of landscape restoration on evapotranspiration and water use in the Yellow River Basin, China. *Acta Geophys.* **72**, 341–356 (2024).
- Xu, S., Yu, Z., Yang, C., Ji, X. & Zhang, K. Trends in evapotranspiration and their responses to climate change and vegetation greening over the upper reaches of the Yellow River Basin. Agric. For. Meteorol. 263, 118–129 (2018).
- Ren, D. et al. Quantifying natural and anthropogenic impacts on streamflow and sediment load reduction in the upper to middle Yellow River Basin. J. Hydrol. Reg. Stud. 53. 101788 (2024).
- Wang, Y., Shao, M., Zhu, Y. & Liu, Z. Impacts of land use and plant characteristics on dried soil layers in different climatic regions on the Loess Plateau of China. Agric. For. Meteorol. 151. 437–448 (2011).
- Liu, Y. et al. Recent anthropogenic curtailing of Yellow river runoff and sediment load is unprecedented over the past 500 v. Proc. Natl Acad. Sci. USA 117. 18251–18257 (2020).
- Wang, Y. et al. Soil moisture decline in China's monsoon loess critical zone: more a result
 of land-use conversion than climate change. Proc. Natl Acad. Sci. USA 121, e2322127121
 (2024).
- Jiao, C., Wang, S., Song, S. & Fu, B. Long-term and seasonal variation of open-surface water bodies in the Yellow River Basin during 1990–2020. Hydrol. Process. 37, e14846 (2023).
- Wang, Y. et al. Anthropogenic drought in the Yellow River Basin: multifaceted and weakening connections between meteorological and hydrological droughts. J. Hydrol. 619, 129273 (2023).
- Wang-Erlandsson, L. et al. A planetary boundary for green water. Nat. Rev. Earth Environ. https://doi.org/10.1038/s43017-022-00287-8 (2022).
- Sang, S., Li, Y., Hou, C., Zi, S. & Lin, H. The interprovincial green water flow in China and its teleconnected effects on the social economy. *Hydrol. Earth Syst. Sci.* 29, 67–84 (2025).
- Jiang, C., Parteli, E. J. R., Xia, Q. & Shao, Y. Evaluation of precipitation reanalysis products for regional hydrological modelling in the Yellow River Basin. *Theor. Appl. Climatol.* 155, 2605–2626 (2024).
- Iqbal, M., Wen, J., Wang, S., Tian, H. & Adnan, M. Variations of precipitation characteristics during the period 1960–2014 in the source region of the Yellow river, China. J. Arid Land 10, 388–401 (2018).
- Ma, Z. et al. Status and ponder of climate and hydrology changes in the Yellow River Basin. Bull. Chin. Acad. Sci. 35, 52–60 (2020).
- Tian, Q. & Yang, S. Regional climatic response to global warming: trends in temperature and precipitation in the Yellow, Yangtze and Pearl river basins since the 1950s. Quat. Int. 440. 1–11 (2017).
- Yang, D., Yang, Y., Gao, G., Jianping, H. & Jiang, E. Water cycle and soil-water coupling processes in the Yellow River Basin. Bull. Natl Nat. Sci. Found. China 35, 544–551 (2021).
- Liang, K., Liu, S., Bai, P. & Nie, R. The Yellow River Basin becomes wetter or drier? The
 case as indicated by mean precipitation and extremes during 1961–2012. Theor. Appl.
 Climatol. 119, 701–722 (2015).
- Zhang, B., Tian, L., Yang, Y. & He, X. Revegetation does not decrease water yield in the Loess Plateau of China. Geophys. Res. Lett. 49, e2022GL098025 (2022).
- Huang, Y., Cai, J., Yin, H. & Cai, M. Correlation of precipitation to temperature variation in the Huanghe river (Yellow River) basin during 1957–2006. J. Hydrol. 372, 1–8 (2009).
- Lu, M., Ma, Z. & Li, M. A review on the changing water cycle of the Yellow River Basin under changes in climate, vegetation, and human water use. *Trans. Atmospheric Sci.* 46, 801–812 (2023)
- Chen, S., Men, B., Pang, J., Bian, Z. & Wang, H. Historical and projected extreme climate changes in the upper Yellow River Basin, China. Sci. Rep. 15, 19061 (2025).
- Li, C., Lu, T., Fu, B., Wang, S. & Holden, J. Sustainable city development challenged by extreme weather in a warming world. Geogr. Sustain. 3, 114–118 (2022).

- Liu, C. & Zheng, H. Trend analysis of hydrological components in the Yellow River Basin. J. Nat. Resour. 18, 129–135 (2003).
- Oki, T. & Kanae, S. Global hydrological cycles and world water resources. Science 313, 1068–1072 (2006).
- Jiang, Z. et al. Revealing the spatio-temporal variability of evapotranspiration and its components based on an improved Shuttleworth-Wallace model in the Yellow River Basin. J. Environ. Manage. 262, 110310 (2020).
- Li, C., Yuan, X., Jiao, Y., Ji, P. & Huang, Z. High-resolution land surface modeling of the irrigation effects on evapotranspiration over the Yellow River Basin. J. Hydrol. 633, 130986 (2024).
- Li, H., Ye, A., Zhang, Y. & Zhao, W. InterComparison and evaluation of multisource soil moisture products in China. Earth Space Sci. 8. e2021EA001845 (2021).
- Lou, D. et al. Changes of soil moisture from multiple sources during 1988–2010 in the Yellow River Basin, China. Adv. Meteorol. 2018, 1950529 (2018).
- Jia, X., Shao, M., Zhu, Y. & Luo, Y. Soil moisture decline due to afforestation across the Loess Plateau, China. J. Hydrol. 546, 113–122 (2017).
- Wang, Q. et al. Will large-scale forestation lead to a soil water deficit crisis in China's drylands? Sci. Bull. 69, 1506–1514 (2024).
- Zhao, Z. et al. Impacts of the grain for green project on soil moisture in the Yellow River Basin, China. Hydrol. Process. 39, e70112 (2025).
- Zhang, K., Xie, X., Zhu, B., Meng, S. & Yao, Y. Unexpected groundwater recovery with decreasing agricultural irrigation in the Yellow River Basin. Agric. Water Manag. 213, 858–867 (2019).
- Lu, M. et al. Lag response of groundwater to changes in water and sediment characteristics in the lower Yellow river, China. J. Hydrol. 612, 128048 (2022).
- Zhang, D., Liu, X., Simmons, C. T., Zhang, L. & Zhang, Q. Changes in groundwater levels across China from 2005 to 2016. J. Hydrol. 623, 129781 (2023).
- Lv, M., Ma, Z., Li, M. & Zheng, Z. Quantitative analysis of terrestrial water storage changes under the grain for green program in the Yellow River Basin. J. Geophys. Res. Atmos. 124, 1336–1351 (2019).
- Zheng, H., Zhang, L., Liu, C., Shao, Q. & Fukushima, Y. Changes in stream flow regime in headwater catchments of the Yellow River Basin since the 1950s. Hydrol. Process. 21, 886–893 (2007).
- Hu, Y., Maskey, S., Uhlenbrook, S. & Zhao, H. Streamflow trends and climate linkages in the source region of the Yellow river, China. Hydrol. Process. 25, 3399–3411 (2011).
- Li, L., Shen, H., Dai, S., Xiao, J. & Shi, X. Response of runoff to climate change and its future tendency in the source region of Yellow river. J. Geogr. Sci. 22, 431–440 (2012).
- Guo, B., Niu, Y., Mantravadi, V. S., Zhang, L. & Liu, G. The variation of rainfall runoff after vegetation restoration in upper reaches of the Yellow river by the remote sensing technology. *Environ. Sci. Pollut. Res.* 28, 50707–50717 (2021).
- Ni, Y. et al. Intra-annual variation in the attribution of runoff evolution in the Yellow river source area. CATENA 225, 107032 (2023).
- Guo, Y. et al. Warming leads to both earlier and later snowmelt floods over the past 70 years. Nat. Commun. 16, 3663 (2025).
- Han, J. et al. Streamflow seasonality in a snow-dwindling world. Nature 629, 1075–1081 (2024).
- Yao, T. et al. The imbalance of the Asian water tower. Nat. Rev. Earth Environ. 3, 618–632 (2022).
- Han, S. et al. Groundwater resource and eco-environmental problem of the Yellow River Basin. Geol. China 48, 1001–1019 (2021).
- Wang, H. & Hu, P. Key issues of ecological conservation in the Yellow River Basin from a water cycle perspective. J. Hydraul. Eng. 51, 1009–1014 (2020).
- Ran, L. & Lu, X. X. Delineation of reservoirs using remote sensing and their storage estimate: an example of the Yellow River Basin, China. Hydrol. Process. 26, 1215–1229 (2012).
- Chen, Y. et al. Monthly monitoring of inundated areas and water storage dynamics in China's large reservoirs using multisource remote sensing. Water Resour. Res. 60, e2023WR036450 (2024).
- Wen, S. et al. Attribution of streamflow changes during 1961–2019 in the upper Yangtze and the upper Yellow River Basins. Clim. Change 177, 60 (2024).
- Wu, X. et al. Evolution and effects of the social-ecological system over a millennium in China's Loess Plateau. Sci. Adv. 6, eabc0276 (2020).
- Luan, J. et al. Estimating hydrological consequences of vegetation greening. J. Hydrol. 611. 128018 (2022).
- Liu, C. et al. Detection of vegetation coverage changes in the Yellow River Basin from 2003 to 2020. Ecol. Indic. 138, 108818 (2022).
- 73. Yang, L. et al. Runoff changes in the major river basins of China and their responses to potential driving forces. *J. Hydrol.* **607**, 127536 (2022).
- Zhang, B., Tian, L., Zhao, X. & Wu, P. Feedbacks between vegetation restoration and local precipitation over the Loess Plateau in China. Sci. China Earth Sci. 64, 920–931 (2021).
- Cao, M. et al. Revegetation impacts on moisture recycling and precipitation trends in the Chinese Loess Plateau. Water Resour. Res. 60, e2024WR038199 (2024).
- Song, S. et al. Sediment transport under increasing anthropogenic stress: regime shifts within the Yellow River, China. Ambio 49, 2015–2025 (2020).
- Ren, M. Sediment discharge of the Yellow River, China: past, present and future a synthesis. Acta Oceanol. Sin. 34, 1–8 (2015).
- Kong, D. et al. The hydro-environmental response on the lower Yellow River to the water-sediment regulation scheme. Ecol. Eng. 79, 69–79 (2015).

- Peng, J., Chen, S. & Dong, P. Temporal variation of sediment load in the Yellow River basin, China, and its impacts on the lower reaches and the river delta. CATENA 83, 135–147 (2010).
- Xu, J. Sedimentation rates in the lower Yellow River over the past 2300 years as influenced by human activities and climate change. *Hydrol. Process.* 17, 3359–3371 (2003)
- Milliman, J. D. & Meade, R. H. World-wide delivery of river sediment to the oceans. J. Geol. 91, 1–21 (1983).
- 82. Syvitski, J. P. M., Vörösmarty, C. J., Kettner, A. J. & Green, P. Impact of humans on the flux of terrestrial sediment to the global coastal ocean. Science **308**, 376–380 (2005).
- Yu, Y. et al. New discharge regime of the Huanghe (Yellow River): causes and implications. Cont. Shelf Res. 69, 62–72 (2013).
- 84. Wang, H. et al. Recent changes in sediment delivery by the Huanghe (Yellow River) to the sea: causes and environmental implications in its estuary. *J. Hydrol.* **391**, 302–313 (2010).
- Pietz, D. A. The Yellow River: the Problem of Water in Modern China (Havard Univ. Press, 2015).
- Chu, Z. The dramatic changes and anthropogenic causes of erosion and deposition in the lower Yellow (Huanghe) River since 1952. Geomorphology 216, 171-179 (2014).
- 87. Yu, S.-Y. et al. Human disturbances dominated the unprecedentedly high frequency of Yellow River flood over the last millennium. Sci. Adv. 9, eadf8576 (2023).
- Chen, Y. Flood dynamics of the lower Yellow River over the last 3000 years: characteristics and implications for geoarchaeology. Quat. Int. 521, 147-157 (2019).
- Jiao, C. et al. River stabilization reshaped human-nature interactions in the lower Yellow River floodplain. J. Environ. Manage. 371, 122957 (2024).
- Xia, J., Li, X., Zhang, X. & Li, T. Recent variation in reach-scale bankfull discharge in the Lower Yellow River. Earth Surf. Process. Landf. 39, 723–734 (2014).
- Kong, D., Latrubesse, E. M., Miao, C. & Zhou, R. Morphological response of the lower Yellow River to the operation of Xiaolangdi Dam, China. Geomorphology 350, 106931 (2020).
- 92. Xu, X., Chen, Z. & Feng, Z. From natural driving to artificial intervention: changes of the Yellow River estuary and delta development. Ocean Coast. Manag. 174, 63–70 (2019).
- Saito, Y., Yang, Z. & Hori, K. The Huanghe (Yellow River) and Changjiang (Yangtze River) deltas: a review on their characteristics, evolution and sediment discharge during the Holocene. Geomorphology 41, 219–231 (2001).
- 94. Cui, B., Yang, Q., Yang, Z. & Zhang, K. Evaluating the ecological performance of wetland restoration in the Yellow River delta, China. Ecol. Eng. 35, 1090–1103 (2009).
- Bi, N. et al. Phase change in evolution of the modern Huanghe (Yellow River) delta: process, pattern, and mechanisms. Mar. Geol. 437, 106516 (2021).
- Wu, X., Feng, X., Fu, B., Yin, S. & He, C. Managing erosion and deposition to stabilize a silt-laden river. Sci. Total Environ. 881, 163444 (2023).
- Li, J., Xia, J. & Ji, Q. Rapid and long-distance channel incision in the Lower Yellow River owing to upstream damming. CATENA 196, 104943 (2021).
- Cheng, Y., Xia, J., Zhou, M., Deng, S. & Li, Z. Adjustment of flood discharge capacity with varying boundary conditions in a braided reach of the lower Yellow River. J. Geogr. Sci. 31, 1615–1632 (2021).
- Miao, C., Kong, D., Wu, J. & Duan, Q. Functional degradation of the water-sediment regulation scheme in the lower Yellow River: spatial and temporal analyses. Sci. Total Environ. 551–552. 16–22 (2016).
- Wu, X. et al. Ecological restoration in the Yellow River Basin enhances hydropower potential. Nat. Commun. 16, 2566 (2025).
- Kong, D. et al. Environmental impact assessments of the Xiaolangdi reservoir on the most hyperconcentrated laden river, Yellow River, China. Environ. Sci. Pollut. Res. 24, 4337–4351 (2017).
- Ma, H. et al. Amplification of downstream flood stage due to damming of fine-grained rivers. Nat. Commun. 13, 3054 (2022).
- Kong, D., Miao, C., Zheng, H. & Gou, J. Dynamic evolution characteristics of the Yellow River delta in response to estuary diversion and a water–sediment regulation scheme. J. Hydrol. 627, 130447 (2023).
- 104. Fu, Y., Chen, S., Ji, H., Fan, Y. & Li, P. The modern Yellow River delta in transition: causes and implications. *Mar. Geol.* **436**, 106476 (2021).
- Wang, H. et al. Impacts of the dam-orientated water-sediment regulation scheme on the lower reaches and delta of the Yellow River (Huanghe): a review. Glob. Planet. Change 157, 93–113 (2017).
- Syvitski, J. P. M. et al. Sinking deltas due to human activities. Nat. Geosci. 2, 681–686 (2009).
- Ren, C. et al. Ageing threatens sustainability of smallholder farming in China. Nature 616, 96–103 (2023).
- Yang, D. et al. How does urbanization process affect ecological landscape pattern?
 An empirical analysis based on scale effects. Ecol. Indic. 154, 110921 (2023).
- Greve, P. et al. Global assessment of water challenges under uncertainty in water scarcity projections. Nat. Sustain. 1, 486–494 (2018).
- Veldkamp, T. I. E. et al. Water scarcity hotspots travel downstream due to human interventions in the 20th and 21st century. Nat. Commun. 8, 15697 (2017).
- Zhang, W., Liang, W., Gao, X., Li, J. & Zhao, X. Trajectory in water scarcity and potential water savings benefits in the Yellow River basin. J. Hydrol. 633, 130998 (2024).
- Yin, Z. et al. Irrigation, damming, and streamflow fluctuations of the Yellow River. Hydrol. Earth Syst. Sci. 25, 1133–1150 (2021).
- 113. Yang, D. et al. Analysis of water resources variability in the Yellow River of China during the last half century using historical data. Water Resour. Res. 40, W06502 (2004).

- 114. Cong, Z., Yang, D., Gao, B., Yang, H. & Hu, H. Hydrological trend analysis in the Yellow River basin using a distributed hydrological model. Water Resour. Res. 45, WOOA13 (2009)
- 115. Singh, D., Xu, M., Singh, N. & Lei, F. Perspectives on emerging pressures and their integrated impact on large river systems: an insight from the Yellow River basin. J. Environ. Manage. 298, 113423 (2021).
- Zhen, N., Rutherfurd, I. & Webber, M. Ecological water, a new focus of China's water management. Sci. Total Environ. 879, 163001 (2023).
- Notice of the Water Allocation Scheme (1998) in the Yellow River http://www.yrcc.gov.cn/ zwzc/zcfq/202312/t20231220 374740.html (Ministry of Water Resources, 2009).
- Liu, Y., Zheng, H. & Zhao, J. Reframing water demand management: a new co-governance framework coupling supply-side and demand-side solutions toward sustainability. Hydrol. Farth Syst. Sci. 28, 2223–2238 (2024).
- Long, D. et al. South-to-north water diversion stabilizing Beijing's groundwater levels.
 Nat. Commun. 11, 3665 (2020)
- Liu, J. et al. Water scarcity assessments in the past, present, and future. Earths Future 5, 545–559 (2017).
- Guo, L., Song, X. & Wang, Q. Simulation of population size and economic scale supportable by the Yellow River's available freshwater in 2030 under multiple scenarios. Humanit. Soc. Sci. Commun. 11, 1–8 (2024).
- 122. Yan, X. et al. Human deforestation outweighed climate as factors affecting Yellow River floods and erosion on the Chinese Loess Plateau since the 10th century. Quat. Sci. Rev. 295, 107796 (2022).
- Bryan, B. A. et al. China's response to a national land-system sustainability emergency. Nature 559, 193–204 (2018).
- Yan, T., Chen, X. & Han, G. A historical examination on land desertification in North China. Chin. Geogr. Sci. 11, 115–119 (2001).
- Qiu, M., Liu, Y., Tian, F., Wang, S. & Fu, B. Marsh decrease was much faster than the water increase among the Yellow River Source wetlands during 1986–2022. Sci. Total Environ. https://doi.org/10.1016/j.scitotenv.2024.174377 (2024).
- 126. Yang, J., Ding, Y. & Chen, R. Spatial and temporal of variations of alpine vegetation cover in the source regions of the Yangtze and Yellow Rivers of the Tibetan Plateau from 1982 to 2001. *Environ. Geol.* **50**, 313–322 (2006).
- Sha, A. et al. Accelerated river meander migration on the Tibetan Plateau caused by permafrost thaw. Geophys. Res. Lett. 52, e2024GL111536 (2025).
- Gao, H., Wang, M. & Hao, X. Check dams in the Yellow River basin: sediment reduction efficiency and future development. *Land Degrad. Dev.* 35, 4042-4054 (2024).
- 129. Xu, X., Zhang, H. & Zhang, O. Development of check-dam systems in gullies on the Loess Plateau, China. Environ. Sci. Policy 7, 79–86 (2004).
- Zhao, G., Mu, X., Wen, Z., Wang, F. & Gao, P. Soil erosion, conservation, and eco-environment changes in the Loess Plateau of China. *Land Degrad. Dev.* 24, 499–510 (2013).
- Soil and Water Conservation Bulletin in Yellow River Basin (Ministry of Water Resources, 2023); http://yrcc.gov.cn/gzfw/stbcgb/index.html.
- Liang, W. et al. Quantifying the impacts of climate change and ecological restoration on streamflow changes based on a Budyko hydrological model in China's Loess Plateau.
 Water Resour. Res. 51, 6500–6519 (2015).
- 133. China's Three-North Shelterbelt Forest Program a Landmark in Desertification Control. The State Council of the People's Republic of China https://english.www.gov.cn/ news/202406/06/content_WS6661298ac6d0868f4e8e7dd0.html (2024).
- 134. Notice of the National Forestry Administration on Issuing the 'Management Measures for the Natural Forest Protection Program' and the 'Verification and Inspection Measures for the Natural Forest Protection Program'. National Forestry Administration http://www.hljcourt.gov.cn/lawdb/show.php?fid=37295 (2001).
- Liu, J., Li, S., Ouyang, Z., Tam, C. & Chen, X. Ecological and socioeconomic effects of China's policies for ecosystem services. Proc. Natl Acad. Sci. USA 105, 9477–9482 (2008).
- Regulations on grain for Green Program. State Council of the People's Republic of China https://www.gov.cn/gongbao/content/2003/content_62531.htm (2002).
- 137. Fu, F. et al. Locating hydrologically unsustainable areas for supporting ecological restoration in China's drylands. *Earths Future* **12**, e2023EF004216 (2024).
- Wang, K. et al. Impacts of vegetation restoration on soil erosion in the Yellow River Basin, China. CATENA 234, 107547 (2024).
- Chen, Q. et al. A review of land use change and its influence in the source region of the Yellow River. Resour. Sci. 42, 446-459 (2020).
- Liu, X. et al. Spatiotemporal changes in vegetation coverage and its driving factors in the three-river headwaters region during 2000–2011. J. Geogr. Sci. 24, 288–302 (2014).
- 141. Chen, Y. et al. Balancing green and grain trade. *Nat. Geosci.* **8**, 739–741 (2015).
- Chen, Y. et al. 65% cover is the sustainable vegetation threshold on the Loess Plateau. Environ. Sci. Ecotechnol. 22, 100442 (2024).
- 143. Ringler, C. et al. Yellow River basin: living with scarcity. Water Int. 35, 681-701 (2010).
- 144. Wang, H. & Ma, T. Optimal water resource allocation considering virtual water trade in the Yellow River Basin. Sci. Rep. 14, 79 (2024).
- 145. Feng, K., Siu, Y. L., Guan, D. & Hubacek, K. Assessing regional virtual water flows and water footprints in the Yellow River Basin, China: a consumption based approach. Appl. Geogr. 32, 691–701 (2012).
- Wei, J. et al. Estimation and influencing factors of agricultural water efficiency in the Yellow River basin, China. J. Clean. Prod. 308, 127249 (2021).

- Wang, J. & Tian, C. The water-carbon constraints' impact on the development of coal power industry in the Yellow River Basin. *Environ. Res. Commun.* 4, 075010 (2022).
- Notice of the Water Allocation Scheme (1987) in the Yellow River. Ministry of Water Resources of the People's Republic of China https://www.gov.cn/zhengce/ zhengceku/2011-03/30/content_3138.htm (1987).
- 149. Lee Kuan Yew Water Prize 2010 Citation for Yellow River Conservancy Commission. Singapore International Water Week https://www.siww.com.sg/landing-page/lee-kuan-yew-water-prize-laureates/yellow-river-conservancy-commission (2010).
- Yang, T. et al. A spatial assessment of hydrologic alteration caused by dam construction in the middle and lower Yellow River, China. Hydrol. Process. 22, 3829–3843 (2008).
- Kong, D. et al. Xiaolangdi Dam: a valve for streamflow extremes on the lower Yellow River. J. Hydrol. 606, 127426 (2022).
- Xia, C. & Pahl-Wostl, C. The development of water allocation management in the Yellow River Basin. Water Resour. Manag. 26, 3395–3414 (2012).
- 153. Wang, H. R., Dong, Y. Y., Wang, Y. & Liu, Q. Water right institution and strategies of the Yellow River Valley. *Water Resour. Manag.* **22**, 1499–1519 (2008).
- 154. Fang, L. & Zhang, L. Does the trading of water rights encourage technology improvement and agricultural water conservation? Agric. Water Manag. 233, 106097 (2020).
- 155. Notice of the Integrated Fertilizers and Water Use management in Agriculture (2016–2020). Ministry of Agriculture of the People's Republic of China https://www.moa.gov.cn/nybgb/2016/diwuqi/201711/t20171127_5920793.htm (2017).
- 156. The South-to-North Water Diversion Project. Ministry of Water Resources of the People's Republic of China http://www.mwr.gov.cn/english/Projects/MagnificentProjects/202311/ t20231102_1690665.html (2019).
- Notice of the Strict Water Management. The State Council of the People's Republic of China http://www.gov.cn/zwgk/2012-02/16/content_2067664.htm (2012).
- Yellow River Protection Law of the People's Republic of China. National People's Congress http://en.npc.gov.cn.cdurl.cn/2022-10/30/c_954870.htm (2022).
- 159. Huggins, X. et al. Hotspots for social and ecological impacts from freshwater stress and storage loss. *Nat. Commun.* **13**, 439 (2022).
- Kwakkel, J. H., Walker, W. E. & Haasnoot, M. Coping with the wickedness of public policy problems: approaches for decision making under deep uncertainty. J. Water Resour. Plan. Manag. 142, 01816001 (2016).
- Wu, W. et al. Beyond engineering: a review of reservoir management through the lens of wickedness, competing objectives and uncertainty. *Environ. Model. Softw.* 167, 105777 (2023).
- 162. Cumming, G. S. & von Cramon-Taubadel, S. Linking economic growth pathways and environmental sustainability by understanding development as alternate social-ecological regimes. Proc. Natl Acad. Sci. USA 115, 9533–9538 (2018).
- Di Baldassarre, G. et al. Water shortages worsened by reservoir effects. Nat. Sustain. 1, 617–622 (2018).
- Gohari, A. et al. Water transfer as a solution to water shortage: a fix that can backfire.
 J. Hydrol. 491, 23–39 (2013).
- 165. Bouckaert, F. W., Wei, Y., Pittock, J., Vasconcelos, V. & Ison, R. River basin governance enabling pathways for sustainable management: a comparative study between Australia, Brazil, China and France. Ambio 51, 1871–1888 (2022).
- Wang, Z. & Zheng, Z. Things and current significance of the Yellow River Water allocation scheme in 1987. Yellow River 41, 109–127 (2019).
- Loch, A., Adamson, D. & Dumbrell, N. P. The fifth stage in water management: policy lessons for water governance. Water Resour. Res. 56, e2019WR026714 (2020).
- Zhang, Y., Yang, P., Liu, X. & Adeloye, A. J. Simulation and optimization coupling model for soil salinization and waterlogging control in the Urad irrigation area, North China. J. Hydrol. 607, 127408 (2022).
- 169. Kandasamy, J. et al. Socio-hydrologic drivers of the pendulum swing between agricultural development and environmental health: a case study from Murrumbidgee River basin, Australia. Hydrol. Earth Syst. Sci. 18, 1027-1041 (2014).
- Ning, T., Li, Z. & Liu, W. Separating the impacts of climate change and land surface alteration on runoff reduction in the Jing River catchment of China. CATENA 147, 80–86 (2016).
- Liu, J., Zhang, Q., Singh, V. P. & Shi, P. Contribution of multiple climatic variables and human activities to streamflow changes across China. J. Hydrol. 545, 145–162 (2017)
- Ge, J., Pitman, A. J., Guo, W., Zan, B. & Fu, C. Impact of revegetation of the Loess Plateau of China on the regional growing season water balance. *Hydrol. Earth Syst. Sci.* 24, 515–533 (2020).
- Wang, X. et al. Gainers and losers of surface and terrestrial water resources in China during 1989–2016. Nat. Commun. 11, 3471 (2020).
- Li, B.-B. et al. Deep soil moisture limits the sustainable vegetation restoration in arid and semi-arid Loess Plateau. Geoderma 399, 115122 (2021).
- Zhao, G. et al. Changing trends and regime shift of streamflow in the Yellow River basin. Stoch. Environ. Res. Risk Assess. 29, 1331–1343 (2015).
- Piao, S. et al. The impacts of climate change on water resources and agriculture in China. Nature 467, 43–51 (2010).
- Moudi, M., Xu, Z., Yao, L. & Yuan, H. Dynamic optimization model for improving Urban water supply system fragility with uncertain streamflow. Water Resour. Manag. 34, 1465–1477 (2020).
- Liu, J. et al. Timing the first emergence and disappearance of global water scarcity.
 Nat. Commun. 15, 7129 (2024).

- 179. Tyulin, A. E., Chursin, A. A., Ragulina, J. V., Akberdina, V. V. & Yudin, A. V. The development of Kondratieff's theory of long waves: the place of the AI economy humanization in the 'competencies-innovations-markets' model. *Humanit*. Soc. Sci. Commun. 10, 1–13 (2023).
- Zhu, Y., Zhang, Y., Zhu, X., Luo, Y. & Tu, Z. The evolution of animal-based dietary structure has contributed to the increase of healthcare expenditures in China. *Humanit. Soc. Sci. Commun.* 11, 1–14 (2024).
- Yin, Y., Tang, Q., Liu, X. & Zhang, X. Water scarcity under various socio-economic pathways and its potential effects on food production in the Yellow River basin. Hydrol. Earth Syst. Sci. 21, 791–804 (2016).
- Te Wierik, S. A. et al. Critical importance of tree and non-tree vegetation for African precipitation. Geophys. Res. Lett. 51, e2023GL103274 (2024).
- Qin, Y., Wang, D., Ziegler, A. D., Fu, B. & Zeng, Z. Impact of Amazonian deforestation on precipitation reverses between seasons. *Nature* 639, 102-108 (2025).
- Zemp, D. C. et al. Self-amplified Amazon forest loss due to vegetation-atmosphere feedbacks. Nat. Commun. 8, 14681 (2017)
- van de Vijsel, R. C., Scheffer, M. & Hoitink, A. J. F. Tipping points in river deltas. Nat. Rev. Earth Environ. https://doi.org/10.1038/s43017-024-00610-5 (2024).
- Binh, D. V. et al. Long-term alterations of flow regimes of the Mekong River and adaptation strategies for the Vietnamese Mekong delta. J. Hydrol. Reg. Stud. 32, 100742 (2020).
- Cox, J. R., Leuven, J. R. F. W., Pierik, H. J., van Egmond, M. & Kleinhans, M. G. Sediment deficit and morphological change of the Rhine–Meuse river mouth attributed to multi-millennial anthropogenic impacts. Cont. Shelf Res. 244, 104766 (2022).
- Dai, Z., Liu, J. T., Wei, W. & Chen, J. Detection of the three Gorges Dam influence on the Changjiang (Yangtze River) submerged delta. Sci. Rep. 4, 6600 (2014).
- 189. Yu, Y. et al. Transboundary cooperation in infrastructure operation generates economic and environmental co-benefits in the Lancang-Mekong River Basin. Nat. Water https:// doi.org/10.1038/s44221-024-00246-1 (2024).
- Grafton, R. Q. et al. Global insights into water resources, climate change and governance. Nat. Clim. Change 3, 315–321 (2013).
- Migliavacca, M. et al. The three major axes of terrestrial ecosystem function. Nature 598, 468–472 (2021)
- Bassani, F., Fatichi, S., Rinaldo, A. & Bonetti, S. Toward a metabolic theory of catchments: scaling of water and carbon fluxes with size. Proc. Natl Acad. Sci. USA 121, e2410736121 (2024)
- 193. Zhao, M. et al. Ecological restoration impact on total terrestrial water storage. Nat. Sustain. 4, 56–62 (2021).
- Koldasbayeva, D. et al. Challenges in data-driven geospatial modeling for environmental research and practice. Nat. Commun. 15, 10700 (2024).
- 195. Razavi, S. et al. Convergent and transdisciplinary integration: on the future of integrated modeling of human-water systems. Water Resour. Res. 61, e2024WR038088 (2025).
- Lubell, M. & Morrison, T. H. Institutional navigation for polycentric sustainability governance. Nat. Sustain. https://doi.org/10.1038/s41893-021-00707-5 (2021).
- Xu, Y. et al. A daily temperature dataset over China and its application in validating a RCM simulation. Adv. Atmospheric Sci. 26, 763–772 (2009).
- Martens, B. et al. GLEAM v3: satellite-based land evaporation and root-zone soil moisture. Geosci. Model Dev. 10, 1903–1925 (2017).
- Hersbach, H. et al. The ERA5 global reanalysis. Q. J. R. Meteorol. Soc. 146, 1999–2049 (2020).
- Tuinenburg, O. A., Theeuwen, J. J. E. & Staal, A. High-resolution global atmospheric moisture connections from evaporation to precipitation. *Earth Syst. Sci. Data* 12, 3177–3188 (2020).
- Wu, J. & Gao, X. A gridded daily observation dataset over China region and comparison with the other datasets. Chin. J. Geophys. 56, 1102–1111 (2013).
- Han, J. et al. A new daily gridded precipitation dataset for the Chinese mainland based on gauge observations. Earth Syst. Sci. Data 15, 3147–3161 (2023).
- He, J. et al. The first high-resolution meteorological forcing dataset for land process studies over China. Sci. Data 7, 25 (2020).
- 204. Miralles, D. G. et al. Global land-surface evaporation estimated from satellite-based observations. Hydrol. Earth Syst. Sci. 15, 453–469 (2011).

- 205. Running, S., Mu, Q. & Zhao, M. MOD16A2 MODIS/Terra net evapotranspiration 8-day l4 global 500m SIN grid V006. NASA Land Processes Distributed Active Archive Center https://doi.org/10.5067/MODIS/MOD16A2.006 (2017).
- Elnashar, A., Wang, L., Wu, B., Zhu, W. & Zeng, H. Synthesis of Global Actual Evapotranspiration from 1982 to 2019. Earth Syst. Sci. Data 13, 447–480 (2021).
- Entekhabi, D. et al. The soil moisture active passive (SMAP) mission. Proc. IEEE 98, 704–716 (2010).
- Rodell, M. et al. The global land data assimilation system. Bull. Am. Meteorol. Soc. 85, 381–394 (2004).
- Kong, D. et al. Evolution of the Yellow River Delta and its relationship with runoff and sediment load from 1983 to 2011. J. Hydrol. 520, 157–167 (2015).
- Chu, Z. X., Sun, X. G., Zhai, S. K. & Xu, K. H. Changing pattern of accretion/erosion of the modern Yellow River (Huanghe) subaerial delta, China: based on remote sensing images. *Mar. Geol.* 227, 13–30 (2006).
- Zhou, F. et al. Deceleration of China's human water use and its key drivers. Proc. Natl Acad. Sci. USA 117, 7702–7711 (2020).
- Klein Goldewijk, K. History database of the global environment 3.3. https://public.yoda. uu.nl/geo/UU01/94FNH0.html (2024).
- 213. Cao, S. et al. Spatiotemporally consistent global dataset of the GIMMS leaf area index (GIMMS LAI4g) from 1982 to 2020. *Earth Syst. Sci. Data* **15**, 4877–4899 (2023).
- Yang, J. & Huang, X. The 30m annual land cover dataset and its dynamics in China from 1990 to 2019. Earth Syst. Sci. Data 13, 3907–3925 (2021).

Acknowledgements

S.W. acknowledges support from the National Natural Science Foundation of China (grant nos. 42041007 and U2243601). S.S. was supported by the National Natural Science Foundation of China (42401126). X.W. was supported by the National Natural Science Foundation of China (42201306). B.F. is grateful for the 111 Project of China (B23027) and the Fundamental Research Funds for the Central Universities. P.R. and S.S. thank the Max Planck Society for funding. J.B. is grateful for support through the Jack and Richard Threet Chair in Sedimentary Geology at Illinois.

Author contributions

S.W., S.S. and B.F. formulated the review and identified the themes to be covered. S.S., H.Z., L.Y. and C.J. drafted the figures and wrote the first draft of the manuscript. P.R. helped structure and revise the manuscript; J.B., X.W., C.L. and W.Z. revised the manuscript and provided extensive comments. All authors made substantial contributions to the discussion of content.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at https://doi.org/10.1038/s43017-025-00718-2.

Peer review information Nature Reviews Earth & Environment thanks Jianhua Ping, Shimin Tian, Jana Cox and the other, anonymous, reviewer for their contribution to the peer review of this work.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© Springer Nature Limited 2025